

Figure 1:

to the cross angular power *only from those objects common to both catalogues*. If it is assumed that in the most extreme case the *IRAS* galaxies are simply a subset of the optical galaxies then the shot noise could be $\simeq 500$. This has a small ($\lesssim 2\%$) effect on the final results when added to the expectation values. The other major approximation here is the assumption of a 4π sky coverage in the statistical comparison below. A full analysis would require the cross-talk of the incomplete sky harmonic coefficients to be taken in to account (Lahav, this volume). This is ignored here, consequently any errors are underestimated.

For a chosen power spectrum a maximum likelihood analysis can be performed for any number of harmonics to determine the best $b_I b_O \sigma_{8,matter}^2$. I have performed this analysis on the optical and *IRAS* catalogues described above for standard (shape parameter $\Gamma = 0.5$) and low density ($\Gamma = 0.2$) CDM models. Using harmonics $l = 1$ to $l = 10$ (to avoid non-linear scales) I find values of $b_I b_O \sigma_{8,matter}^2 = 0.82 \pm 0.05$ for standard CDM and $b_I b_O \sigma_{8,matter}^2 = 0.84 \pm 0.06$ for low density CDM. The joint normalisation is therefore remarkably indifferent to the assumed power spectrum (at least on these scales of 8 to $50h^{-1} Mpc$). From other studies [3],[1],[4] the real space value of $b_I \sigma_{8,matter}$ has been determined as $\simeq 0.7$. It is therefore possible to make a simple estimate of b_O/b_I since $b_O/b_I = (b_I b_O \sigma_{8,matter}^2)/(b_I \sigma_{8,matter})^2$. I therefore estimate $b_O/b_I \simeq 1.7$ on the scales of 8 to $50h^{-1} Mpc$ from this analysis. As a cautionary note, Figure 1 shows the harmonic reconstruction number density as a function of distance (in arbitrary units) along a line close to the SuperGalactic Plane (using harmonics to $l_{max} = 10$) for the optical (solid curve) and *IRAS* (dashed curve) galaxies. The reconstructions have been renormalised according to the mean density of optical and *IRAS* galaxies. Centaurus/Great Attractor and Perseus-Pisces are labelled, the vertical dotted lines are at $b = -90^\circ, 0^\circ$ and 90° . Clearly, while both populations trace the same underlying structure there is a marked difference at the peaks (the differences in sample depths is also a factor). Work in progress will address the issue of relative bias as a function of scale, using the SHA approach.


References

- [1] Fisher K. B., Davis M., Strauss M. A., Yahil A., Huchra J. P., 1993. *Astrophys. J.*, 402,42
- [2] Hudson M. J., 1993. *Mon. Not. R. astr. Soc.*, in press
- [3] Lahav O., Nemiroff R. J., Piran T., 1990. *Astrophys. J.*, 350, 119
- [4] Scharf C. A., Hoffman Y., Lahav O., Lynden-Bell D., 1992. *Mon. Not. R. astr. Soc.*, 256, 229
- [5] Strauss M. A., Davis M., Yahil A., Huchra J. P., 1990. *Astrophys. J.*, 385,421
- [6] Yahil A., Strauss M. A., Davis M., Huchra J. P., 1991. *Astrophys. J.*, 372, 380

COSMOLOGICAL PARAMETERS FROM SPHERICAL HARMONIC ANALYSES

Caleb. A. Scharf

Institute of Astronomy, Madingley Rd., Cambridge, CB3 0HA.



Abstract

In the rest of this volume Lahav and Fisher have presented the method of Spherical Harmonic Analysis (SHA) and its use in estimating cosmologically interesting parameters. Here I demonstrate the use of SHA in cross-correlating galaxy catalogues and present an estimate of the ratio of the optical and *IRAS* galaxy bias parameters; $b_O/b_I \simeq 1.7$.

1 METHOD AND RESULTS

The spherical harmonic transform of the distribution of objects on the full sky can be found as a sum over objects $a_{lm} = \sum_i f(r_i) Y_{lm}^*(\hat{\mathbf{r}}_i)$, where f is an arbitrary radial weight and Y_{lm} are the orthonormal spherical harmonics. An estimator of the cross angular power between two catalogues (I and O) can be written as:

$$c_{l,IO} = \frac{1}{(\Omega_{obs}/4\pi)(2l+1)} [Re(a_{l0,I})Re(a_{l0,O}) + 2 \sum_{m=1}^l (Re(a_{lm,I})Re(a_{lm,O}) + Im(a_{lm,I})Im(a_{lm,O}))] . \quad (1)$$

The model prediction can be written as:

$$\langle a_{l,I} a_{l,O} \rangle = b_I b_O \sigma_{8,matter}^2 \frac{2}{\pi} \int k^2 P(k) \Psi_{l,I}^R(k) \Psi_{l,O}^R(k) dk , \quad (2)$$

where $b_I b_O$ is the product of the *IRAS* and optical bias parameters, $\sigma_{8,matter}^2$ is the rms variance in density in $8h^{-1} Mpc$ spheres, $P(k)$ is the unnormalised power spectrum of matter fluctuations and $\Psi_{l,I}^R$ and $\Psi_{l,O}^R$ are the appropriate harmonic window functions for *IRAS* and optical galaxies in real space, as described by Lahav (this volume). Here a simple number weighted ($f(r) = 1$) harmonic cross-correlation is applied to the *2Jy IRAS* catalogue [5] and a merged UGC and ESO optical catalogue constructed by M. Hudson [2]. The optical catalogue has a sky coverage of $\simeq 67\%$ (12177 objects), here I have used a *2Jy* catalogue with unsurveyed regions filled according to [6] and have then imposed the same limited coverage as in the optical (leaving 2137 objects). There will be a shot noise contribution